

## **POSTOPERATIVE INSTRUCTIONS FOR BROWLIFT**

Our goal is for you to achieve the best possible result from your surgery. This requires participation from both you and our staff during the postoperative period. You should be familiar with these instructions in order to make the postoperative course a smooth one. We are happy to answer any questions or concerns you may have regarding your postoperative care.

- **SWELLING:** Keeping the head elevated will help with swelling.
- **WHEN TO CALL THE DOCTOR:** If any signs of infection such as an area of redness, oral temperature over 101F, foul smelling drainage, or pain which is not relieved with medication, then you must call immediately to be evaluated. Dr. Peters' office number is **(985) 223-2602**. After 5 PM, weekends, and holidays, your call will be transferred to our answering service.
- **ACTIVITY:** Avoid any strenuous physical activity including running and lifting children. Keep any lifting under 10 pounds. Most of the discomfort is during the first 2 weeks following surgery. You will not begin to feel "back to normal" for about 3-4 weeks. You may begin to go on leisurely walks after 2 weeks. **DO NOT OVER DO IT.** Too much activity will prolong the healing time.
- **STAPLES:** Your incisions are closed with staples, which will be removed in about 14-18 days.
- **EXPOSURE TO SUNLIGHT:** It is recommended that you protect the incision lines from the sun for at least one year following surgery. Application of sunscreen is recommended under a bathing suit or clothes, since sunlight can reach the skin.
- **SMOKING:** Advise that you do not use any tobacco products for at least 14 days following your surgery.
- **ALCOHOL:** Alcohol dilates the blood vessels and may increase the chance of post-operative bleeding. It also contributes to dehydration. Please do not drink until you have stopped taking the prescription pain pills as this combination can be dangerous.
- **DRIVING:** Please do not drive for at least 10 days following your surgery. You should not drive while taking the prescription pain medication
- **ITCHING:** (without hives): Benadryl 25-50 mg every 6 hours
- **GAS:** (caused by pain medication): Phazyme gel tabs.
- **CONSTIPATION:** (Caused by pain medication): Fruit juices, eat canned fruit, or Colace or Senecot stool softeners.
- **FEVER:** (without chills or body aches and less than 100 degrees): Two Tylenol every four hours. Look carefully at you incisions for any possible signs of infection.
- **POSTOPERATIVE APPOINTMENTS:** It is very important that you follow the scheduled appointments we establish for you after surgery.

You may continue to use your prescribed narcotic medication as directed. Please ask Dr. Peters prior to initiating any other medications such as Ibuprofen or Vitamin E.

**Mark H. Peters, MD, FACS**

---

I have reviewed the postoperative care instructions with \_\_\_\_\_; I understand these instructions and a copy has been provided to me.

Patient \_\_\_\_\_ Date \_\_\_\_\_