

**POSTOPERATIVE INSTRUCTIONS CHIN AUGMENTATION
And HOW TO PREPARE**

Our goal is for you to achieve the best possible result from your surgery. This requires participation from both you and our staff during the postoperative period. You should be familiar with these instructions in order to make the postoperative course a smooth one. We are happy to answer any questions or concerns you may have regarding your postoperative care.

- Keeping the head elevated will help with swelling.
- Mouthwash to rinse the mouth. It may be difficult to brush your teeth at first.
- Safe skincare products which to wash your face for the first time. Make sure all your skincare products are hypo-allergenic and formulated for sensitive and dry skin. (examples: Lubriderm, Eucerin)
- We want you to avoid straining. A stool softener or mild laxative may be needed. (Colace or Senecot)
- Limit your intake with citrus juices. Juices are highly acidic and may sour your stomach.
- Flexible straws for drinking liquids easily in a reclining position.
- A hand-held shower head in the tub or shower enclosure plus a plastic chair upon which to sit. This is a safe way to shower for the first time.
- Leave your splint dressing in place, until your first post op visit.
- Clean visible suture lines with half strength hydrogen peroxide, i.e. ½ water and ½ peroxide. “Dab” rather than wipe.” Do this once or twice daily.
- Lots of patience!!!!!!!!!!!!

GENERAL INFORMATION

- **ACTIVITY:** Most of the discomfort is during the first 2 weeks following surgery. You will not begin to feel “back to normal” for about 3-4 weeks. You may begin to go on leisurely walks after 2 weeks. **DO NOT OVER DO IT.** Too much activity will prolong the healing time.
- **WHEN TO CALL THE DOCTOR:** If any signs of infection such as an area of redness, oral temperature over 101F, foul smelling drainage, or pain which is not relieved with medication, then you must call immediately to be evaluated. Dr. Peters’ office number is **(985) 223-2602**. After 5 PM, weekends, and holidays, your call will be transferred to our answering service.
- **SUTURES:** The stitches in your mouth are dissolvable.
- **SMOKING:** We advise that you do not use any tobacco products for at least 14 days following your surgery.

Mark H. Peters, MD, FACS

- **ALCOHOL:** Alcohol dilates the blood vessels and may increase the chance of post-operative bleeding. It also contributes to dehydration. Please do not drink until you have stopped taking the prescription pain pills as this combination can be dangerous.
- **DRIVING:** Please do not drive for at least 5 days following the surgery. You should not drive while taking the prescription pain medication.
- **ITCHING:** Benadryl (over the counter): 25-50 mg every 6 hours.
- **GAS:** (caused by pain medication): Phazyme gel tabs.
- **CONSTIPATION:** (Caused by pain medication): Fruit juices, eat canned fruit, Colace or Senecot stool softeners.
- **FEVER:** (without chills or body aches and less than 100 degrees): Two Tylenol every four hours. Look carefully at you incisions for any possible signs of infection.
- **POSTOPERATIVE APPOINTMENTS:** It is very important that you follow the scheduled appointments we establish for you after surgery.

You may continue to use your prescribed narcotic medication as directed. Please ask Dr. Peters prior to initiating any other medications such as Ibuprofen or Vitamin E.

I have reviewed the postoperative care instructions with _____; I understand these instructions and a copy has been provided to me.

Patient _____ Date _____